


Class 11 - Chapter 19 Arithmetic Progression



RD Sharma Solutions for Class 11 Maths Chapter 19–Arithmetic Progressions

Class 11: Maths Chapter 19 solutions. Complete Class 11 Maths Chapter 19 Notes.

RD Sharma Solutions for Class 11 Maths Chapter 19–Arithmetic Progressions

RD Sharma 11th Maths Chapter 19, Class 11 Maths Chapter 19 solutions

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>



EXERCISE 19.1 PAGE NO: 19.4

1. If the n^{th} term of a sequence is given by $a_n = n^2 - n + 1$, write down its first five terms.

Solution:

Given:

$$a_n = n^2 - n + 1$$

By using the values $n = 1, 2, 3, 4, 5$ we can find the first five terms.

When $n = 1$:

$$a_1 = (1)^2 - 1 + 1$$

$$= 1 - 1 + 1$$

$$= 1$$

When $n = 2$:

$$a_2 = (2)^2 - 2 + 1$$

$$= 4 - 2 + 1$$

$$= 3$$

When $n = 3$:

$$a_3 = (3)^2 - 3 + 1$$

$$= 9 - 3 + 1$$

$$= 7$$

When $n = 4$:

$$a_4 = (4)^2 - 4 + 1$$

$$= 16 - 4 + 1$$

$$= 13$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

When $n = 5$:

$$a_5 = (5)^2 - 5 + 1$$

$$= 25 - 5 + 1$$

$$= 21$$

\therefore First five terms of the sequence are 1, 3, 7, 13, 21.

2. A sequence is defined by $a_n = n^3 - 6n^2 + 11n - 6$, $n \in \mathbf{N}$. Show that the first three terms of the sequence are zero and all other terms are positive.

Solution:

Given:

$$a_n = n^3 - 6n^2 + 11n - 6, n \in \mathbf{N}$$

By using the values $n = 1, 2, 3$ we can find the first three terms.

When $n = 1$:

$$a_1 = (1)^3 - 6(1)^2 + 11(1) - 6$$

$$= 1 - 6 + 11 - 6$$

$$= 12 - 12$$

$$= 0$$

When $n = 2$:

$$a_2 = (2)^3 - 6(2)^2 + 11(2) - 6$$

$$= 8 - 6(4) + 22 - 6$$

$$= 8 - 24 + 22 - 6$$

$$= 30 - 30$$

$$= 0$$

When $n = 3$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$\begin{aligned}a_3 &= (3)^3 - 6(3)^2 + 11(3) - 6 \\&= 27 - 6(9) + 33 - 6 \\&= 27 - 54 + 33 - 6 \\&= 60 - 60 \\&= 0\end{aligned}$$

This shows that the first three terms of the sequence is zero.

Now, let's check for when $n = n$:

$$\begin{aligned}a_n &= n^3 - 6n^2 + 11n - 6 \\&= n^3 - 6n^2 + 11n - 6 - n + n - 2 + 2 \\&= n^3 - 6n^2 + 12n - 8 - n + 2 \\&= (n)^3 - 3 \times 2n(n - 2) - (2)^3 - n + 2\end{aligned}$$

By using the formula, $\{(a - b)^3 = (a)^3 - (b)^3 - 3ab(a - b)\}$

$$a_n = (n - 2)^3 - (n - 2)$$

Here, $n - 2$ will always be positive for $n > 3$

$\therefore a_n$ is always positive for $n > 3$

3. Find the first four terms of the sequence defined by $a_1 = 3$ and $a_n = 3a_{n-1} + 2$, for all $n > 1$.

Solution:

Given:

$$a_1 = 3 \text{ and } a_n = 3a_{n-1} + 2, \text{ for all } n > 1$$

By using the values $n = 1, 2, 3, 4$ we can find the first four terms.

When $n = 1$:

$$a_1 = 3$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

When $n = 2$:

$$a_2 = 3a_{2-1} + 2$$

$$= 3a_1 + 2$$

$$= 3(3) + 2$$

$$= 9 + 2$$

$$= 11$$

When $n = 3$:

$$a_3 = 3a_{3-1} + 2$$

$$= 3a_2 + 2$$

$$= 3(11) + 2$$

$$= 33 + 2$$

$$= 35$$

When $n = 4$:

$$a_4 = 3a_{4-1} + 2$$

$$= 3a_3 + 2$$

$$= 3(35) + 2$$

$$= 105 + 2$$

$$= 107$$

\therefore First four terms of sequence are 3, 11, 35, 107.

4. Write the first five terms in each of the following sequences:

(i) $a_1 = 1, a_n = a_{n-1} + 2, n > 1$

(ii) $a_1 = 1 = a_2, a_n = a_{n-1} + a_{n-2}, n > 2$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

(iii) $a_1 = a_2 = 2, a_n = a_{n-1} - 1, n > 2$

Solution:

(i) $a_1 = 1, a_n = a_{n-1} + 2, n > 1$

By using the values $n = 1, 2, 3, 4, 5$ we can find the first five terms.

Given:

$$a_1 = 1$$

When $n = 2$:

$$a_2 = a_{2-1} + 2$$

$$= a_1 + 2$$

$$= 1 + 2$$

$$= 3$$

When $n = 3$:

$$a_3 = a_{3-1} + 2$$

$$= a_2 + 2$$

$$= 3 + 2$$

$$= 5$$

When $n = 4$:

$$a_4 = a_{4-1} + 2$$

$$= a_3 + 2$$

$$= 5 + 2$$

$$= 7$$

When $n = 5$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a_5 = a_{5-1} + 2$$

$$= a_4 + 2$$

$$= 7 + 2$$

$$= 9$$

∴ First five terms of the sequence are 1, 3, 5, 7, 9.

$$(ii) a_1 = 1 = a_2, a_n = a_{n-1} + a_{n-2}, n > 2$$

By using the values $n = 1, 2, 3, 4, 5$ we can find the first five terms.

Given:

$$a_1 = 1$$

$$a_2 = 1$$

When $n = 3$:

$$a_3 = a_{3-1} + a_{3-2}$$

$$= a_2 + a_1$$

$$= 1 + 1$$

$$= 2$$

When $n = 4$:

$$a_4 = a_{4-1} + a_{4-2}$$

$$= a_3 + a_2$$

$$= 2 + 1$$

$$= 3$$

When $n = 5$:

$$a_5 = a_{5-1} + a_{5-2}$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= a_4 + a_3$$

$$= 3 + 2$$

$$= 5$$

∴ First five terms of the sequence are 1, 1, 2, 3, 5.

(iii) $a_1 = a_2 = 2$, $a_n = a_{n-1} - 1$, $n > 2$

By using the values $n = 1, 2, 3, 4, 5$ we can find the first five terms.

Given:

$$a_1 = 2$$

$$a_2 = 2$$

When $n = 3$:

$$a_3 = a_{3-1} - 1$$

$$= a_2 - 1$$

$$= 2 - 1$$

$$= 1$$

When $n = 4$:

$$a_4 = a_{4-1} - 1$$

$$= a_3 - 1$$

$$= 1 - 1$$

$$= 0$$

When $n = 5$:

$$a_5 = a_{5-1} - 1$$

$$= a_4 - 1$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 0 - 1$$

$$= -1$$

∴ First five terms of the sequence are 2, 2, 1, 0, -1.

5. The Fibonacci sequence is defined by $a_1 = 1 = a_2$, $a_n = a_{n-1} + a_{n-2}$ for $n > 2$. Find $(a_{n+1})/a_n$ for $n = 1, 2, 3, 4, 5$.

Solution:

Given:

$$a_1 = 1$$

$$a_2 = 1$$

$$a_n = a_{n-1} + a_{n-2}$$

When $n = 1$:

$$(a_{n+1})/a_n = (a_{1+1})/a_1$$

$$= a_2/a_1$$

$$= 1/1$$

$$= 1$$

$$a_3 = a_{3-1} + a_{3-2}$$

$$= a_2 + a_1$$

$$= 1 + 1$$

$$= 2$$

When $n = 2$:

$$(a_{n+1})/a_n = (a_{2+1})/a_2$$

$$= a_3/a_2$$

$$= 2/1$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 2$$

$$a_4 = a_{4-1} + a_{4-2}$$

$$= a_3 + a_2$$

$$= 2 + 1$$

$$= 3$$

When $n = 3$:

$$(a_{n+1})/a_n = (a_{3+1})/a_3$$

$$= a_4/a_3$$

$$= 3/2$$

$$a_5 = a_{5-1} + a_{5-2}$$

$$= a_4 + a_3$$

$$= 3 + 2$$

$$= 5$$

When $n = 4$:

$$(a_{n+1})/a_n = (a_{4+1})/a_4$$

$$= a_5/a_4$$

$$= 5/3$$

$$a_6 = a_{6-1} + a_{6-2}$$

$$= a_5 + a_4$$

$$= 5 + 3$$

$$= 8$$

When $n = 5$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$(a_{n+1})/a_n = (a_{5+1})/a_5$$

$$= a_6/a_5 = 8/5$$

∴ Value of $(a_{n+1})/a_n$ when $n = 1, 2, 3, 4, 5$ are $1, 2, 3/2, 5/3, 8/5$

EXERCISE 19.2 PAGE NO: 19.11

1. Find:

(i) 10th term of the A.P. 1, 4, 7, 10,

(ii) 18th term of the A.P. $\sqrt{2}, 3\sqrt{2}, 5\sqrt{2}, \dots$

(iii) nth term of the A.P 13, 8, 3, -2,

Solution:

(i) 10th term of the A.P. 1, 4, 7, 10,

Arithmetic Progression (AP) whose common difference is $= a_n - a_{n-1}$ where $n > 0$

Let us consider, $a = a_1 = 1, a_2 = 4 \dots$

So, Common difference, $d = a_2 - a_1 = 4 - 1 = 3$

To find the 10th term of A.P, firstly find a_n

By using the formula,

$$a_n = a + (n-1) d$$

$$= 1 + (n-1) 3$$

$$= 1 + 3n - 3$$

$$= 3n - 2$$

When $n = 10$:

$$a_{10} = 3(10) - 2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 30 - 2$$

$$= 28$$

Hence, 10th term is 28.

(ii) 18th term of the A.P. $\sqrt{2}, 3\sqrt{2}, 5\sqrt{2}, \dots$

Arithmetic Progression (AP) whose common difference is $= a_n - a_{n-1}$ where $n > 0$

Let us consider, $a = a_1 = \sqrt{2}, a_2 = 3\sqrt{2} \dots$

So, Common difference, $d = a_2 - a_1 = 3\sqrt{2} - \sqrt{2} = 2\sqrt{2}$

To find the 18th term of A.P, firstly find a_n

By using the formula,

$$a_n = a + (n-1) d$$

$$= \sqrt{2} + (n - 1) 2\sqrt{2}$$

$$= \sqrt{2} + 2\sqrt{2}n - 2\sqrt{2}$$

$$= 2\sqrt{2}n - \sqrt{2}$$

When $n = 18$:

$$a_{18} = 2\sqrt{2}(18) - \sqrt{2}$$

$$= 36\sqrt{2} - \sqrt{2}$$

$$= 35\sqrt{2}$$

Hence, 10th term is $35\sqrt{2}$

(iii) nth term of the A.P 13, 8, 3, -2,

Arithmetic Progression (AP) whose common difference is $= a_n - a_{n-1}$ where $n > 0$

Let us consider, $a = a_1 = 13, a_2 = 8 \dots$

So, Common difference, $d = a_2 - a_1 = 8 - 13 = -5$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

To find the n^{th} term of A.P, firstly find a_n

By using the formula,

$$\begin{aligned}a_n &= a + (n-1) d \\&= 13 + (n-1) (-5) \\&= 13 - 5n + 5 \\&= 18 - 5n\end{aligned}$$

Hence, n^{th} term is $18 - 5n$

2. In an A.P., show that $a_{m+n} + a_{m-n} = 2a_m$.

Solution:

We know the first term is 'a' and the common difference of an A.P is d.

Given:

$$a_{m+n} + a_{m-n} = 2a_m$$

By using the formula,

$$a_n = a + (n - 1)d$$

Now, let us take LHS: $a_{m+n} + a_{m-n}$

$$\begin{aligned}a_{m+n} + a_{m-n} &= a + (m + n - 1)d + a + (m - n - 1)d \\&= a + md + nd - d + a + md - nd - d \\&= 2a + 2md - 2d \\&= 2(a + md - d) \\&= 2[a + d(m - 1)] \{ \because a_n = a + (n - 1)d \}\end{aligned}$$

$$a_{m+n} + a_{m-n} = 2a_m$$

Hence Proved.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

3. (i) Which term of the A.P. 3, 8, 13,... is 248 ?

(ii) Which term of the A.P. 84, 80, 76,... is 0 ?

(iii) Which term of the A.P. 4, 9, 14,... is 254 ?

Solution:

(i) Which term of the A.P. 3, 8, 13,... is 248 ?

Given A.P is 3, 8, 13,...

Here, $a_1 = a = 3$, $a_2 = 8$

Common difference, $d = a_2 - a_1 = 8 - 3 = 5$

We know, $a_n = a + (n - 1)d$

$$a_n = 3 + (n - 1)5$$

$$= 3 + 5n - 5$$

$$= 5n - 2$$

Now, to find which term of A.P is 248

Put $a_n = 248$

$$\therefore 5n - 2 = 248$$

$$= 248 + 2$$

$$= 250$$

$$= 250/5$$

$$= 50$$

Hence, 50th term of given A.P is 248.

(ii) Which term of the A.P. 84, 80, 76,... is 0 ?

Given A.P is 84, 80, 76,...

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Here, $a_1 = a = 84$, $a_2 = 88$

Common difference, $d = a_2 - a_1 = 88 - 84 = 4$

We know, $a_n = a + (n - 1)d$

$$a_n = 84 + (n - 1) \cdot 4$$

$$= 84 + 4n - 4$$

$$= 80 + 4n$$

Now, to find which term of A.P is 0

Put $a_n = 0$

$$80 + 4n = 0$$

$$4n = -80$$

$$n = -80/4$$

$$= -20$$

Hence, 20th term of given A.P is 0.

(iii) Which term of the A.P. 4, 9, 14,... is 254 ?

Given A.P is 4, 9, 14,...

Here, $a_1 = a = 4$, $a_2 = 9$

Common difference, $d = a_2 - a_1 = 9 - 4 = 5$

We know, $a_n = a + (n - 1)d$

$$a_n = 4 + (n - 1) \cdot 5$$

$$= 4 + 5n - 5$$

$$= 5n - 1$$

Now, to find which term of A.P is 254

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$\text{Put } a_n = 254$$

$$5n - 1 = 254$$

$$5n = 254 + 1$$

$$5n = 255$$

$$n = 255/5$$

$$= 51$$

Hence, 51st term of given A.P is 254.

4. (i) Is 68 a term of the A.P. 7, 10, 13,...?

(ii) Is 302 a term of the A.P. 3, 8, 13,...?

Solution:

(i) Is 68 a term of the A.P. 7, 10, 13,...?

Given A.P is 7, 10, 13,...

Here, $a_1 = a = 7$, $a_2 = 10$

Common difference, $d = a_2 - a_1 = 10 - 7 = 3$

We know, $a_n = a + (n - 1)d$ [where, a is first term or a_1 and d is common difference and n is any natural number]

$$a_n = 7 + (n - 1)3$$

$$= 7 + 3n - 3$$

$$= 3n + 4$$

Now, to find whether 68 is a term of this A.P. or not

$$\text{Put } a_n = 68$$

$$3n + 4 = 68$$

$$3n = 68 - 4$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$3n = 64$$

$$n = 64/3$$

64/3 is not a natural number

Hence, 68 is not a term of given A.P.

(ii) Is 302 a term of the A.P. 3, 8, 13, ...?

Given A.P is 3, 8, 13, ...

$$\text{Here, } a_1 = a = 3, a_2 = 8$$

$$\text{Common difference, } d = a_2 - a_1 = 8 - 3 = 5$$

We know, $a_n = a + (n - 1)d$

$$a_n = 3 + (n - 1)5$$

$$= 3 + 5n - 5$$

$$= 5n - 2$$

To find whether 302 is a term of this A.P. or not

$$\text{Put } a_n = 302$$

$$5n - 2 = 302$$

$$5n = 302 + 2$$

$$5n = 304$$

$$n = 304/5$$

304/5 is not a natural number

Hence, 304 is not a term of given A.P.

5. (i) Which term of the sequence $24, 23 \frac{1}{4}, 22 \frac{1}{2}, 21 \frac{3}{4}$ is the first negative term?

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Given:

AP: $24, 23 \frac{1}{4}, 22 \frac{1}{2}, 21 \frac{3}{4}, \dots = 24, 93/4, 45/2, 87/4, \dots$

Here, $a_1 = a = 24, a_2 = 93/4$

Common difference, $d = a_2 - a_1 = 93/4 - 24$

$$= (93 - 96)/4$$

$$= -3/4$$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

We know, $a_n = a + (n - 1) d$

$$a_n = 24 + (n - 1) (-3/4)$$

$$= 24 - 3/4n + 3/4$$

$$= (96+3)/4 - 3/4n$$

$$= 99/4 - 3/4n$$

Now we need to find, first negative term.

Put $a_n < 0$

$$a_n = 99/4 - 3/4n < 0$$

$$99/4 < 3/4n$$

$$3n > 99$$

$$n > 99/3$$

$$n > 33$$

Hence, 34th term is the first negative term of given AP.

(ii) Which term of the sequence $12 + 8i, 11 + 6i, 10 + 4i, \dots$ is (a) purely real (b) purely imaginary ?

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Solution:

Given:

AP: $12 + 8i, 11 + 6i, 10 + 4i, \dots$

Here, $a_1 = a = 12 + 8i, a_2 = 11 + 6i$

Common difference, $d = a_2 - a_1$

$$= 11 + 6i - (12 + 8i)$$

$$= 11 - 12 + 6i - 8i$$

$$= -1 - 2i$$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

$$a_n = 12 + 8i + (n - 1) (-1 - 2i)$$

$$= 12 + 8i - n - 2ni + 1 + 2i$$

$$= 13 + 10i - n - 2ni$$

$$= (13 - n) + (10 - 2n) i$$

To find purely real term of this A.P., imaginary part have to be zero

$$10 - 2n = 0$$

$$2n = 10$$

$$n = 10/2$$

$$= 5$$

Hence, 5th term is purely real.

To find purely imaginary term of this A.P., real part have to be zero

$$\therefore 13 - n = 0$$

$$n = 13$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Hence, 13th term is purely imaginary.

6. (i) How many terms are in A.P. 7, 10, 13,...43?

Solution:

Given:

AP: 7, 10, 13,...

Here, $a_1 = a = 7$, $a_2 = 10$

Common difference, $d = a_2 - a_1 = 10 - 7 = 3$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

$$a_n = 7 + (n - 1)3$$

$$= 7 + 3n - 3$$

$$= 3n + 4$$

To find total terms of the A.P., put $a_n = 43$ as 43 is last term of A.P.

$$3n + 4 = 43$$

$$3n = 43 - 4$$

$$3n = 39$$

$$n = 39/3$$

$$= 13$$

Hence, total 13 terms exists in the given A.P.

(ii) How many terms are there in the A.P. -1, -5/6, -2/3, -1/2, ..., 10/3 ?

Solution:

Given:

AP: -1, -5/6, -2/3, -1/2, ...

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Here, $a_1 = a = -1$, $a_2 = -5/6$

Common difference, $d = a_2 - a_1$

$$= -5/6 - (-1)$$

$$= -5/6 + 1$$

$$= (-5+6)/6$$

$$= 1/6$$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

$$a_n = -1 + (n - 1) 1/6$$

$$= -1 + 1/6n - 1/6$$

$$= (-6-1)/6 + 1/6n$$

$$= -7/6 + 1/6n$$

To find total terms of the AP,

Put $a_n = 10/3$ [Since, $10/3$ is the last term of AP]

$$a_n = -7/6 + 1/6n = 10/3$$

$$1/6n = 10/3 + 7/6$$

$$1/6n = (20+7)/6$$

$$1/6n = 27/6$$

$$n = 27$$

Hence, total 27 terms exists in the given A.P.

7. The first term of an A.P. is 5, the common difference is 3, and the last term is 80; find the number of terms.

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Given:

First term, $a = 5$; last term, $l = a_n = 80$

Common difference, $d = 3$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

$$a_n = 5 + (n - 1)3$$

$$= 5 + 3n - 3$$

$$= 3n + 2$$

To find total terms of the A.P., put $a_n = 80$ as 80 is last term of A.P.

$$3n + 2 = 80$$

$$3n = 80 - 2$$

$$3n = 78$$

$$n = 78/3$$

$$= 26$$

Hence, total 26 terms exists in the given A.P.

8. The 6th and 17th terms of an A.P. are 19 and 41 respectively. Find the 40th term.

Solution:

Given:

6th term of an A.P is 19 and 17th terms of an A.P. is 41

So, $a_6 = 19$ and $a_{17} = 41$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

When $n = 6$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a_6 = a + (6 - 1) d$$

$$= a + 5d$$

Similarly, When $n = 17$:

$$a_{17} = a + (17 - 1)d$$

$$= a + 16d$$

According to question:

$$a_6 = 19 \text{ and } a_{17} = 41$$

$$a + 5d = 19 \text{ (i)}$$

$$\text{And } a + 16d = 41 \text{ (ii)}$$

Let us subtract equation (i) from (ii) we get,

$$a + 16d - (a + 5d) = 41 - 19$$

$$a + 16d - a - 5d = 22$$

$$11d = 22$$

$$d = 22/11$$

$$= 2$$

put the value of d in equation (i):

$$a + 5(2) = 19$$

$$a + 10 = 19$$

$$a = 19 - 10$$

$$= 9$$

$$\text{As, } a_n = a + (n - 1)d$$

$$a_{40} = a + (40 - 1)d$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= a + 39d$$

Now put the value of $a = 9$ and $d = 2$ in a_{40} we get,

$$a_{40} = 9 + 39(2)$$

$$= 9 + 78$$

$$= 87$$

Hence, 40th term of the given A.P. is 87.

9. If 9th term of an A.P. is Zero, prove that its 29th term is double the 19th term.

Solution:

Given:

9th term of an A.P is 0

$$\text{So, } a_9 = 0$$

We need to prove: $a_{29} = 2a_{19}$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is common difference and n is any natural number]

When $n = 9$:

$$a_9 = a + (9 - 1)d$$

$$= a + 8d$$

According to question:

$$a_9 = 0$$

$$a + 8d = 0$$

$$a = -8d$$

When $n = 19$:

$$a_{19} = a + (19 - 1)d$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= a + 18d$$

$$= -8d + 18d$$

$$= 10d$$

When $n = 29$:

$$a_{29} = a + (29 - 1)d$$

$$= a + 28d$$

$$= -8d + 28d \text{ [Since, } a = -8d]$$

$$= 20d$$

$$= 2 \times 10d$$

$$a_{29} = 2a_{19} \text{ [Since, } a_{19} = 10d]$$

Hence Proved.

10. If 10 times the 10th term of an A.P. is equal to 15 times the 15th term, show that the 25th term of the A.P. is Zero.

Solution:

Given:

10 times the 10th term of an A.P. is equal to 15 times the 15th term

$$\text{So, } 10a_{10} = 15a_{15}$$

We need to prove: $a_{25} = 0$

We know, $a_n = a + (n - 1)d$ [where a is first term or a_1 and d is common difference and n is any natural number]

When $n = 10$:

$$a_{10} = a + (10 - 1)d$$

$$= a + 9d$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

When $n = 15$:

$$a_{15} = a + (15 - 1)d$$

$$= a + 14d$$

When $n = 25$:

$$a_{25} = a + (25 - 1)d$$

$$= a + 24d \dots\dots\dots(i)$$

According to question:

$$10a_{10} = 15a_{15}$$

$$10(a + 9d) = 15(a + 14d)$$

$$10a + 90d = 15a + 210d$$

$$10a - 15a + 90d - 210d = 0$$

$$-5a - 120d = 0$$

$$-5(a + 24d) = 0$$

$$a + 24d = 0$$

$$a_{25} = 0 \text{ [From (i)]}$$

Hence Proved.

11. The 10th and 18th term of an A.P. are 41 and 73 respectively, find 26th term.

Solution:

Given:

10th term of an A.P is 41, and 18th terms of an A.P. is 73

So, $a_{10} = 41$ and $a_{18} = 73$

We know, $a_n = a + (n - 1) d$ [where a is first term or a_1 and d is the common difference and n is any natural number]

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

When $n = 10$:

$$a_{10} = a + (10 - 1)d$$

$$= a + 9d$$

When $n = 18$:

$$a_{18} = a + (18 - 1)d$$

$$= a + 17d$$

According to question:

$$a_{10} = 41 \text{ and } a_{18} = 73$$

$$a + 9d = 41 \text{(i)}$$

$$\text{And } a + 17d = 73 \text{(ii)}$$

Let us subtract equation (i) from (ii) we get,

$$a + 17d - (a + 9d) = 73 - 41$$

$$a + 17d - a - 9d = 32$$

$$8d = 32$$

$$d = 32/8$$

$$d = 4$$

Put the value of d in equation (i) we get,

$$a + 9(4) = 41$$

$$a + 36 = 41$$

$$a = 41 - 36$$

$$a = 5$$

we know, $a_n = a + (n - 1)d$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a_{26} = a + (26 - 1)d$$

$$= a + 25d$$

Now put the value of $a = 5$ and $d = 4$ in a_{26}

$$a_{26} = 5 + 25(4)$$

$$= 5 + 100$$

$$= 105$$

Hence, 26th term of the given A.P. is 105.

12. In a certain A.P. the 24th term is twice the 10th term. Prove that the 72nd term is twice the 34th term.

Solution:

Given:

24th term is twice the 10th term

$$\text{So, } a_{24} = 2a_{10}$$

We need to prove: $a_{72} = 2a_{34}$

We know, $a_n = a + (n - 1)d$ [where a is first term or a_1 and d is common difference and n is any natural number]

When $n = 10$:

$$a_{10} = a + (10 - 1)d$$

$$= a + 9d$$

When $n = 24$:

$$a_{24} = a + (24 - 1)d$$

$$= a + 23d$$

When $n = 34$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a_{34} = a + (34 - 1)d$$
$$= a + 33d \dots\dots(i)$$

When $n = 72$:

$$a_{72} = a + (72 - 1)d$$
$$= a + 71d$$

According to question:

$$a_{24} = 2a_{10}$$

$$a + 23d = 2(a + 9d)$$

$$a + 23d = 2a + 18d$$

$$a - 2a + 23d - 18d = 0$$

$$-a + 5d = 0$$

$$a = 5d$$

Now, $a_{72} = a + 71d$

$$a_{72} = 5d + 71d$$

$$= 76d$$

$$= 10d + 66d$$

$$= 2(5d + 33d)$$

$$= 2(a + 33d) \text{ [since, } a = 5d]$$

$$a_{72} = 2a_{34} \text{ (From (i))}$$

Hence Proved.

EXERCISE 19.3 PAGE NO: 19.15

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

1. The Sum of the three terms of an A.P. is 21 and the product of the first, and the third terms exceed the second term by 6, find three terms.

Solution:

Given:

The sum of first three terms is 21

Let us assume the first three terms as $a - d$, a , $a + d$ [where a is the first term and d is the common difference]

So, sum of first three terms is

$$a - d + a + a + d = 21$$

$$3a = 21$$

$$a = 7$$

It is also given that product of first and third term exceeds the second by 6

$$\text{So, } (a - d)(a + d) - a = 6$$

$$a^2 - d^2 - a = 6$$

Substituting the value of $a = 7$, we get

$$7^2 - d^2 - 7 = 6$$

$$d^2 = 36$$

$$d = 6 \text{ or } d = -6$$

Hence, the terms of AP are $a - d$, a , $a + d$ which is 1, 7, 13.

2. Three numbers are in A.P. If the sum of these numbers be 27 and the product 648, find the numbers

Solution:

Given:

Sum of first three terms is 27

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Let us assume the first three terms as $a - d$, a , $a + d$ [where a is the first term and d is the common difference]

So, sum of first three terms is

$$a - d + a + a + d = 27$$

$$3a = 27$$

$$a = 9$$

It is given that the product of three terms is 648

$$\text{So, } a^3 - ad^2 = 648$$

Substituting the value of $a = 9$, we get

$$9^3 - 9d^2 = 648$$

$$729 - 9d^2 = 648$$

$$81 = 9d^2$$

$$d = 3 \text{ or } d = -3$$

Hence, the given terms are $a - d$, a , $a + d$ which is 6, 9, 12.

3. Find the four numbers in A.P., whose sum is 50 and in which the greatest number is 4 times the least.

Solution:

Given:

Sum of four terms is 50.

Let us assume these four terms as $a - 3d$, $a - d$, $a + d$, $a + 3d$

It is given that, sum of these terms is $4a = 50$

$$\text{So, } a = 50/4$$

$$= 25/2 \dots (i)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

It is also given that the greatest number is 4 time the least

$$a + 3d = 4(a - 3d)$$

Substitute the value of $a = 25/2$, we get

$$(25+6d)/2 = 50 - 12d$$

$$30d = 75$$

$$d = 75/30$$

$$= 25/10$$

$$= 5/2 \dots (ii)$$

Hence, the terms of AP are $a - 3d$, $a - d$, $a + d$, $a + 3d$ which is 5, 10, 15, 20

4. The sum of three numbers in A.P. is 12, and the sum of their cubes is 288. Find the numbers.

Solution:

Given:

The sum of three numbers is 12

Let us assume the numbers in AP are $a - d$, a , $a + d$

So,

$$3a = 12$$

$$a = 4$$

It is also given that the sum of their cube is 288

$$(a - d)^3 + a^3 + (a + d)^3 = 288$$

$$a^3 - d^3 - 3ad(a - d) + a^3 + a^3 + d^3 + 3ad(a + d) = 288$$

Substitute the value of $a = 4$, we get

$$64 - d^3 - 12d(4 - d) + 64 + 64 + d^3 + 12d(4 + d) = 288$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$192 + 24d^2 = 288$$

$$d = 2 \text{ or } d = -2$$

Hence, the numbers are $a - d$, a , $a + d$ which is 2, 4, 6 or 6, 4, 2

5. If the sum of three numbers in A.P. is 24 and their product is 440, find the numbers.

Solution:

Given:

Sum of first three terms is 24

Let us assume the first three terms are $a - d$, a , $a + d$ [where a is the first term and d is the common difference]

So, sum of first three terms is $a - d + a + a + d = 24$

$$3a = 24$$

$$a = 8$$

It is given that the product of three terms is 440

$$\text{So } a^3 - ad^2 = 440$$

Substitute the value of $a = 8$, we get

$$8^3 - 8d^2 = 440$$

$$512 - 8d^2 = 440$$

$$72 = 8d^2$$

$$d = 3 \text{ or } d = -3$$

Hence, the given terms are $a - d$, a , $a + d$ which is 5, 8, 11

6. The angles of a quadrilateral are in A.P. whose common difference is 10. Find the angles

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Given: $d = 10$

We know that the sum of all angles in a quadrilateral is 360

Let us assume the angles are $a - 3d$, $a - d$, $a + d$, $a + 3d$

So, $a - 2d + a - d + a + d + a + 2d = 360$

$4a = 360$

$a = 90 \dots$ (i)

And,

$(a - d) - (a - 3d) = 10$

$2d = 10$

$d = 10/2$

$= 5$

Hence, the angles are $a - 3d$, $a - d$, $a + d$, $a + 3d$ which is 75° , 85° , 95° , 105°

EXERCISE 19.4 PAGE NO: 19.30

1. Find the sum of the following arithmetic progressions:

(i) 50, 46, 42, to 10 terms

(ii) 1, 3, 5, 7, ... to 12 terms

(iii) 3, $9/2$, 6, $15/2$, ... to 25 terms

(iv) 41, 36, 31, ... to 12 terms

(v) $a+b$, $a-b$, $a-3b$, ... to 22 terms

(vi) $(x - y)^2$, $(x^2 + y^2)$, $(x + y)^2$, ... to n terms

(vii) $(x - y)/(x + y)$, $(3x - 2y)/(x + y)$, $(5x - 3y)/(x + y)$, ... to n terms

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Solution:

(i) 50, 46, 42, to 10 terms

$$n = 10$$

$$\text{First term, } a = a_1 = 50$$

$$\text{Common difference, } d = a_2 - a_1 = 46 - 50 = -4$$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = 10/2 (100 + (9) (-4))$$

$$= 5 (100 - 36)$$

$$= 5 (64)$$

$$= 320$$

∴ The sum of the given AP is 320.

(ii) 1, 3, 5, 7, ... to 12 terms

$$n = 12$$

$$\text{First term, } a = a_1 = 1$$

$$\text{Common difference, } d = a_2 - a_1 = 3 - 1 = 2$$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = 12/2 (2(1) + (12-1) (2))$$

$$= 6 (2 + (11) (2))$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 6 (2 + 22)$$

$$= 6 (24)$$

$$= 144$$

∴ The sum of the given AP is 144.

(iii) 3, $9/2$, 6, $15/2$, ... to 25 terms

$$n = 25$$

First term, $a = a_1 = 3$

Common difference, $d = a_2 - a_1 = 9/2 - 3 = (9 - 6)/2 = 3/2$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = 25/2 (2(3) + (25-1) (3/2))$$

$$= 25/2 (6 + (24) (3/2))$$

$$= 25/2 (6 + 36)$$

$$= 25/2 (42)$$

$$= 25 (21)$$

$$= 525$$

∴ The sum of the given AP is 525.

(iv) 41, 36, 31, ... to 12 terms

$$n = 12$$

First term, $a = a_1 = 41$

Common difference, $d = a_2 - a_1 = 36 - 41 = -5$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = 12/2 (2(41) + (12-1) (-5))$$

$$= 6 (82 + (11) (-5))$$

$$= 6 (82 - 55)$$

$$= 6 (27)$$

$$= 162$$

∴ The sum of the given AP is 162.

(v) $a+b, a-b, a-3b, \dots$ to 22 terms

$$n = 22$$

First term, $a = a_1 = a+b$

Common difference, $d = a_2 - a_1 = (a-b) - (a+b) = a-b-a-b = -2b$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = 22/2 (2(a+b) + (22-1) (-2b))$$

$$= 11 (2a + 2b + (21) (-2b))$$

$$= 11 (2a + 2b - 42b)$$

$$= 11 (2a - 40b)$$

$$= 22a - 440b$$

∴ The sum of the given AP is $22a - 440b$.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

(vi) $(x - y)^2, (x^2 + y^2), (x + y)^2, \dots$ to n terms

$n = n$

First term, $a = a_1 = (x-y)^2$

Common difference, $d = a_2 - a_1 = (x^2 + y^2) - (x-y)^2 = 2xy$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = n/2 (2(x-y)^2 + (n-1) (2xy))$$

$$= n/2 (2 (x^2 + y^2 - 2xy) + 2xyn - 2xy)$$

$$= n/2 \times 2 ((x^2 + y^2 - 2xy) + xyn - xy)$$

$$= n (x^2 + y^2 - 3xy + xyn)$$

\therefore The sum of the given AP is $n (x^2 + y^2 - 3xy + xyn)$.

(vii) $(x - y)/(x + y), (3x - 2y)/(x + y), (5x - 3y)/(x + y), \dots$ to n terms

$n = n$

First term, $a = a_1 = (x-y)/(x+y)$

Common difference, $d = a_2 - a_1 = (3x - 2y)/(x + y) - (x-y)/(x+y) = (2x - y)/(x+y)$

By using the formula,

$$S = n/2 (2a + (n - 1) d)$$

Substitute the values of 'a' and 'd', we get

$$S = n/2 (2((x-y)/(x+y)) + (n-1) ((2x - y)/(x+y)))$$

$$= n/2(x+y) \{n (2x-y) - y\}$$

\therefore The sum of the given AP is $n/2(x+y) \{n (2x-y) - y\}$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

2. Find the sum of the following series:

(i) $2 + 5 + 8 + \dots + 182$

(ii) $101 + 99 + 97 + \dots + 47$

(iii) $(a - b)^2 + (a^2 + b^2) + (a + b)^2 + \dots + [(a + b)^2 + 6ab]$

Solution:

(i) $2 + 5 + 8 + \dots + 182$

First term, $a = a_1 = 2$

Common difference, $d = a_2 - a_1 = 5 - 2 = 3$

a_n term of given AP is 182

$$a_n = a + (n-1)d$$

$$182 = 2 + (n-1)3$$

$$182 = 2 + 3n - 3$$

$$182 = 3n - 1$$

$$3n = 182 + 1$$

$$n = 183/3$$

$$= 61$$

Now,

By using the formula,

$$S = n/2 (a + l)$$

$$= 61/2 (2 + 182)$$

$$= 61/2 (184)$$

$$= 61 (92)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 5612$$

∴ The sum of the series is 5612

(ii) $101 + 99 + 97 + \dots + 47$

First term, $a = a_1 = 101$

Common difference, $d = a_2 - a_1 = 99 - 101 = -2$

a_n term of given AP is 47

$$a_n = a + (n-1)d$$

$$47 = 101 + (n-1)(-2)$$

$$47 = 101 - 2n + 2$$

$$2n = 103 - 47$$

$$2n = 56$$

$$n = 56/2 = 28$$

Then,

$$S = n/2 (a + l)$$

$$= 28/2 (101 + 47)$$

$$= 28/2 (148)$$

$$= 14 (148)$$

$$= 2072$$

∴ The sum of the series is 2072

(iii) $(a - b)^2 + (a^2 + b^2) + (a + b)^2 + \dots + [(a + b)^2 + 6ab]$

First term, $a = a_1 = (a-b)^2$

Common difference, $d = a_2 - a_1 = (a^2 + b^2) - (a - b)^2 = 2ab$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

a_n term of given AP is $[(a + b)^2 + 6ab]$

$$a_n = a + (n-1)d[(a + b)^2 + 6ab] = (a-b)^2 + (n-1)2ab$$

$$a^2 + b^2 + 2ab + 6ab = a^2 + b^2 - 2ab + 2abn - 2ab$$

$$a^2 + b^2 + 8ab - a^2 - b^2 + 2ab + 2ab = 2abn$$

$$12ab = 2abn$$

$$n = 12ab / 2ab$$

$$= 6$$

Then,

$$S = n/2 (a + l)$$

$$= 6/2 ((a-b)^2 + [(a + b)^2 + 6ab])$$

$$= 3 (a^2 + b^2 - 2ab + a^2 + b^2 + 2ab + 6ab)$$

$$= 3 (2a^2 + 2b^2 + 6ab)$$

$$= 3 \times 2 (a^2 + b^2 + 3ab)$$

$$= 6 (a^2 + b^2 + 3ab)$$

\therefore The sum of the series is $6 (a^2 + b^2 + 3ab)$

3. Find the sum of first n natural numbers.

Solution:

Let AP be 1, 2, 3, 4, ..., n

Here,

First term, $a = a_1 = 1$

Common difference, $d = a_2 - a_1 = 2 - 1 = 1$

$l = n$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

So, the sum of n terms = $S = n/2 [2a + (n-1) d]$

$$= n/2 [2(1) + (n-1) 1]$$

$$= n/2 [2 + n - 1]$$

$$= n/2 [n - 1]$$

\therefore The sum of the first n natural numbers is $n(n-1)/2$

4. Find the sum of all – natural numbers between 1 and 100, which are divisible by 2 or 5

Solution:

The natural numbers which are divisible by 2 or 5 are:

$$2 + 4 + 5 + 6 + 8 + 10 + \dots + 100 = (2 + 4 + 6 + \dots + 100) + (5 + 15 + 25 + \dots + 95)$$

Now, $(2 + 4 + 6 + \dots + 100) + (5 + 15 + 25 + \dots + 95)$ are AP with common difference of 2 and 10.

So, for the 1st sequence $\Rightarrow (2 + 4 + 6 + \dots + 100)$

$$a = 2, d = 4-2 = 2, a_n = 100$$

By using the formula,

$$a_n = a + (n-1)d$$

$$100 = 2 + (n-1)2$$

$$100 = 2 + 2n - 2$$

$$2n = 100$$

$$n = 100/2$$

$$= 50$$

So now, $S = n/2 (2a + (n-1)d)$

$$= 50/2 (2(2) + (50-1)2)$$

$$= 25 (4 + 49(2))$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 25 (4 + 98)$$

$$= 2550$$

Again, for the 2nd sequence, (5 + 15 + 25 + ... + 95)

$$a = 5, d = 15 - 5 = 10, a_n = 95$$

By using the formula,

$$a_n = a + (n-1)d$$

$$95 = 5 + (n-1)10$$

$$95 = 5 + 10n - 10$$

$$10n = 95 + 10 - 5$$

$$10n = 100$$

$$n = 100/10$$

$$= 10$$

$$\text{So now, } S = n/2 (2a + (n-1)d)$$

$$= 10/2 (2(5) + (10-1)10)$$

$$= 5 (10 + 9(10))$$

$$= 5 (10 + 90)$$

$$= 500$$

∴ The sum of the numbers divisible by 2 or 5 is: 2550 + 500 = 3050

5. Find the sum of first n odd natural numbers.

Solution:

Given an AP of first n odd natural numbers whose first term a is 1, and common difference d is 3

The sequence is 1, 3, 5, 7,n

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a = 1, d = 3-1 = 2, n = n$$

By using the formula,

$$S = n/2 [2a + (n-1)d]$$

$$= n/2 [2(1) + (n-1)2]$$

$$= n/2 [2 + 2n - 2]$$

$$= n/2 [2n]$$

$$= n^2$$

∴ The sum of the first n odd natural numbers is n^2 .

6. Find the sum of all odd numbers between 100 and 200

Solution:

The series is 101, 103, 105, ..., 199

Let the number of terms be n

$$\text{So, } a = 101, d = 103 - 101 = 2, a_n = 199$$

$$a_n = a + (n-1)d$$

$$199 = 101 + (n-1)2$$

$$199 = 101 + 2n - 2$$

$$2n = 199 - 101 + 2$$

$$2n = 100$$

$$n = 100/2$$

$$= 50$$

By using the formula,

$$\text{The sum of n terms} = S = n/2[a + l]$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 50/2 [101 + 199]$$

$$= 25 [300]$$

$$= 7500$$

∴ The sum of the odd numbers between 100 and 200 is 7500.

7. Show that the sum of all odd integers between 1 and 1000 which are divisible by 3 is 83667

Solution:

The odd numbers between 1 and 1000 divisible by 3 are 3, 9, 15, ..., 999

Let the number of terms be 'n', so the nth term is 999

$$a = 3, d = 9 - 3 = 6, a_n = 999$$

$$a_n = a + (n-1)d$$

$$999 = 3 + (n-1)6$$

$$999 = 3 + 6n - 6$$

$$6n = 999 + 6 - 3$$

$$6n = 1002$$

$$n = 1002/6$$

$$= 167$$

By using the formula,

$$\text{Sum of } n \text{ terms, } S = n/2 [a + l]$$

$$= 167/2 [3 + 999]$$

$$= 167/2 [1002]$$

$$= 167 [501]$$

$$= 83667$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

∴ The sum of all odd integers between 1 and 1000 which are divisible by 3 is 83667.

Hence proved.

8. Find the sum of all integers between 84 and 719, which are multiples of 5

Solution:

The series is 85, 90, 95, ..., 715

Let there be 'n' terms in the AP

So, $a = 85$, $d = 90 - 85 = 5$, $a_n = 715$

$$a_n = a + (n-1)d$$

$$715 = 85 + (n-1)5$$

$$715 = 85 + 5n - 5$$

$$5n = 715 - 85 + 5$$

$$5n = 635$$

$$n = 635/5$$

$$= 127$$

By using the formula,

$$\text{Sum of } n \text{ terms, } S = n/2 [a + l]$$

$$= 127/2 [85 + 715]$$

$$= 127/2 [800]$$

$$= 127 [400]$$

$$= 50800$$

∴ The sum of all integers between 84 and 719, which are multiples of 5 is 50800.

9. Find the sum of all integers between 50 and 500 which are divisible by 7

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Solution:

The series of integers divisible by 7 between 50 and 500 are 56, 63, 70, ..., 497

Let the number of terms be 'n'

So, $a = 56$, $d = 63 - 56 = 7$, $a_n = 497$

$$a_n = a + (n-1)d$$

$$497 = 56 + (n-1)7$$

$$497 = 56 + 7n - 7$$

$$7n = 497 - 56 + 7$$

$$7n = 448$$

$$n = 448/7$$

$$= 64$$

By using the formula,

Sum of n terms, $S = n/2 [a + l]$

$$= 64/2 [56 + 497]$$

$$= 32 [553]$$

$$= 17696$$

∴ The sum of all integers between 50 and 500 which are divisible by 7 is 17696.

10. Find the sum of all even integers between 101 and 999**Solution:**

We know that all even integers will have a common difference of 2.

So, AP is 102, 104, 106, ..., 998

We know, $a = 102$, $d = 104 - 102 = 2$, $a_n = 998$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

By using the formula,

$$a_n = a + (n-1)d$$

$$998 = 102 + (n-1)2$$

$$998 = 102 + 2n - 2$$

$$2n = 998 - 102 + 2$$

$$2n = 898$$

$$n = 898/2$$

$$= 449$$

By using the formula,

$$\text{Sum of } n \text{ terms, } S = n/2 [a + l]$$

$$= 449/2 [102 + 998]$$

$$= 449/2 [1100]$$

$$= 449 [550]$$

$$= 246950$$

∴ The sum of all even integers between 101 and 999 is 246950.

EXERCISE 19.5 PAGE NO: 19.42

1. If $1/a$, $1/b$, $1/c$ are in A.P., prove that:

(i) $(b+c)/a$, $(c+a)/b$, $(a+b)/c$ are in A.P.

(ii) $a(b+c)$, $b(c+a)$, $c(a+b)$ are in A.P.

Solution:

(i) $(b+c)/a$, $(c+a)/b$, $(a+b)/c$ are in A.P.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

We know that, if a, b, c are in AP, then $b - a = c - b$

If, $1/a, 1/b, 1/c$ are in AP

Then, $1/b - 1/a = 1/c - 1/b$

If $(b+c)/a, (c+a)/b, (a+b)/c$ are in AP

Then, $(c+a)/b - (b+c)/a = (a+b)/c - (c+a)/b$

Let us take LCM

$$\frac{ca + a^2 - b^2 - cb}{ab} = \frac{ab + b^2 - c^2 - ac}{bc}$$

Now let us consider LHS:

$$\frac{ca + a^2 - b^2 - cb}{ab}$$

Multiply both numerator and denominator by 'c', we get,

$$\begin{aligned} \frac{ca + a^2 - b^2 - cb}{ab} &= \frac{c^2a + ca^2 - cb^2 - c^2b}{abc} \\ &= \frac{C(b-a)(a+b+c)}{abc} \end{aligned}$$

Now let us consider RHS:

$$\frac{ab + b^2 - c^2 - ac}{bc}$$

Multiply both numerator and denominator by 'a', we get,

$$\begin{aligned} \frac{ab + b^2 - c^2 - ac}{bc} &= \frac{a^2b + ab^2 - ac^2 - a^2c}{abc} \\ &= \frac{a(b-c)(a+b+c)}{abc} \end{aligned}$$

LHS = RHS

$$\frac{C(b-a)(a+b+c)}{abc} = \frac{a(b-c)(a+b+c)}{abc}$$

Since, $1/a, 1/b, 1/c$ are in AP

$$1/b - 1/a = 1/c - 1/b$$

$$C(b-a) = a(b-c)$$

Hence, the given terms are in AP.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Since, $1/a, 1/b, 1/c$ are in AP

$$1/b - 1/a = 1/c - 1/b$$

$$C(b - a) = a(b - c)$$

Hence, the given terms are in AP.

(ii) $a(b + c), b(c + a), c(a + b)$ are in A.P.

We know that if, $b(c + a) - a(b + c) = c(a + b) - b(c + a)$

Consider LHS:

$$b(c + a) - a(b + c)$$

Upon simplification we get,

$$b(c + a) - a(b + c) = bc + ba - ab - ac$$

$$= c(b - a)$$

Now,

$$c(a + b) - b(c + a) = ca + cb - bc - ba$$

$$= a(c - b)$$

We know,

$1/a, 1/b, 1/c$ are in AP

$$\text{So, } 1/a - 1/b = 1/b - 1/c$$

$$\text{Or } c(b - a) = a(c - b)$$

Hence, given terms are in AP.

2. If a^2, b^2, c^2 are in AP., prove that $a/(b+c), b/(c+a), c/(a+b)$ are in AP.

Solution:

If a^2, b^2, c^2 are in AP then, $b^2 - a^2 = c^2 - b^2$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

If $a/(b+c)$, $b/(c+a)$, $c/(a+b)$ are in AP then,

$$b/(c+a) - a/(b+c) = c/(a+b) - b/(c+a)$$

Let us take LCM on both the sides we get,

$$\frac{b^2 + bc - a^2 - ac}{(a+c)(b+c)} = \frac{ca + c^2 - b^2 - ab}{(a+b)(b+c)}$$

$$\frac{(b-a)(a+b+c)}{(a+c)(b+c)} = \frac{(c-b)(a+b+c)}{(a+b)(b+c)}$$

Since, $b^2 - a^2 = c^2 - b^2$

Substituting $b^2 - a^2 = c^2 - b^2$ in above, we get

LHS = RHS

Hence, given terms are in AP

3. If a, b, c are in A.P., then show that:

(i) $a^2(b+c)$, $b^2(c+a)$, $c^2(a+b)$ are also in A.P.

(ii) $b+c-a$, $c+a-b$, $a+b-c$ are in A.P.

(iii) $bc - a^2$, $ca - b^2$, $ab - c^2$ are in A.P.

Solution:

(i) $a^2(b+c)$, $b^2(c+a)$, $c^2(a+b)$ are also in A.P.

$$\text{If } b^2(c+a) - a^2(b+c) = c^2(a+b) - b^2(c+a)$$

$$b^2c + b^2a - a^2b - a^2c = c^2a + c^2b - b^2a - b^2c$$

$$\text{Given, } b - a = c - b$$

And since a, b, c are in AP,

$$c(b^2 - a^2) + ab(b - a) = a(c^2 - b^2) + bc(c - b)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$(b - a)(ab + bc + ca) = (c - b)(ab + bc + ca)$$

Upon cancelling, $ab + bc + ca$ from both sides

$$b - a = c - b$$

$$2b = c + a \text{ [which is true]}$$

Hence, given terms are in AP

(ii) $b + c - a$, $c + a - b$, $a + b - c$ are in A.P.

$$\text{If } (c + a - b) - (b + c - a) = (a + b - c) - (c + a - b)$$

Then, $b + c - a$, $c + a - b$, $a + b - c$ are in A.P.

Let us consider LHS and RHS

$$(c + a - b) - (b + c - a) = (a + b - c) - (c + a - b)$$

$$2a - 2b = 2b - 2c$$

$$b - a = c - b$$

And since a , b , c are in AP,

$$b - a = c - b$$

Hence, given terms are in AP.

(iii) $bc - a^2$, $ca - b^2$, $ab - c^2$ are in A.P.

$$\text{If } (ca - b^2) - (bc - a^2) = (ab - c^2) - (ca - b^2)$$

Then, $bc - a^2$, $ca - b^2$, $ab - c^2$ are in A.P.

Let us consider LHS and RHS

$$(ca - b^2) - (bc - a^2) = (ab - c^2) - (ca - b^2)$$

$$(a - b^2 - bc + a^2) = (ab - c^2 - ca + b^2)$$

$$(a - b)(a + b + c) = (b - c)(a + b + c)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$a - b = b - c$$

And since a, b, c are in AP,

$$b - c = a - b$$

Hence, given terms are in AP

4. If $(b+c)/a, (c+a)/b, (a+b)/c$ are in AP., prove that:

(i) $1/a, 1/b, 1/c$ are in AP

(ii) bc, ca, ab are in AP

Solution:

(i) $1/a, 1/b, 1/c$ are in AP

If $1/a, 1/b, 1/c$ are in AP then,

$$1/b - 1/a = 1/c - 1/b$$

Let us consider LHS:

$$1/b - 1/a = (a-b)/ab$$

$$= c(a-b)/abc \text{ [by multiplying with 'c' on both the numerator and denominator]}$$

Let us consider RHS:

$$1/c - 1/b = (b-c)/bc$$

$$= a(b-c)/bc \text{ [by multiplying with 'a' on both the numerator and denominator]}$$

Since, $(b+c)/a, (c+a)/b, (a+b)/c$ are in AP

$$\frac{c+a}{b} - \frac{b+c}{a} = \frac{a+b}{c} - \frac{c+a}{b}$$
$$\frac{b^2 + bc - a^2 - ac}{ab} = \frac{ca + c^2 - b^2 - ab}{bc}$$
$$\frac{(b-a)(a+b+c)}{ab} = \frac{(c-b)(a+b+c)}{bc}$$

$$\frac{a(b-c)}{abc} = \frac{c(a-b)}{abc}$$

$$a(b-c) = c(a-b)$$

LHS = RHS

Hence, the given terms are in AP

(ii) bc, ca, ab are in AP

If bc, ca, ab are in AP then,

$$ca - bc = ab - ca$$

$$c(a-b) = a(b-c)$$

If $1/a, 1/b, 1/c$ are in AP then,

$$1/b - 1/a = 1/c - 1/b$$

$$c(a-b) = a(b-c)$$

Hence, the given terms are in AP

5. If a, b, c are in A.P., prove that:

(i) $(a - c)^2 = 4(a - b)(b - c)$

(ii) $a^2 + c^2 + 4ac = 2(ab + bc + ca)$

(iii) $a^3 + c^3 + 6abc = 8b^3$

Solution:

(i) $(a - c)^2 = 4(a - b)(b - c)$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Let us expand the above expression

$$a^2 + c^2 - 2ac = 4(ab - ac - b^2 + bc)$$

$$a^2 + 4c^2b^2 + 2ac - 4ab - 4bc = 0$$

$$(a + c - 2b)^2 = 0$$

$$a + c - 2b = 0$$

Since a, b, c are in AP

$$b - a = c - b$$

$$a + c - 2b = 0$$

$$a + c = 2b$$

Hence, $(a - c)^2 = 4(a - b)(b - c)$

(ii) $a^2 + c^2 + 4ac = 2(ab + bc + ca)$

Let us expand the above expression

$$a^2 + c^2 + 4ac = 2(ab + bc + ca)$$

$$a^2 + c^2 + 2ac - 2ab - 2bc = 0$$

$$(a + c - b)^2 - b^2 = 0$$

$$a + c - b = b$$

$$a + c - 2b = 0$$

$$2b = a + c$$

$$b = (a + c)/2$$

Since a, b, c are in AP

$$b - a = c - b$$

$$b = (a + c)/2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Hence, $a^2 + c^2 + 4ac = 2(ab + bc + ca)$

(iii) $a^3 + c^3 + 6abc = 8b^3$

Let us expand the above expression

$$a^3 + c^3 + 6abc = 8b^3$$

$$a^3 + c^3 - (2b)^3 + 6abc = 0$$

$$a^3 + (-2b)^3 + c^3 + 3a(-2b)c = 0$$

Since, if $a + b + c = 0$, $a^3 + b^3 + c^3 = 3abc$

$$(a - 2b + c)^3 = 0$$

$$a - 2b + c = 0$$

$$a + c = 2b$$

$$b = (a+c)/2$$

Since a, b, c are in AP

$$a - b = c - b$$

$$b = (a+c)/2$$

Hence, $a^3 + c^3 + 6abc = 8b^3$

6. If $a(1/b + 1/c)$, $b(1/c + 1/a)$, $c(1/a + 1/b)$ are in AP., prove that a, b, c are in AP.

Solution:

Here, we know $a(1/b + 1/c)$, $b(1/c + 1/a)$, $c(1/a + 1/b)$ are in AP

Also, $a(1/b + 1/c) + 1$, $b(1/c + 1/a) + 1$, $c(1/a + 1/b) + 1$ are in AP

Let us take LCM for each expression then we get,

$(ac+ab+bc)/bc$, $(ab+bc+ac)/ac$, $(cb+ac+ab)/ab$ are in AP

$1/bc$, $1/ac$, $1/ab$ are in AP

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Let us multiply numerator with 'abc', we get

$abc/bc, abc/ac, abc/ab$ are in AP

$\therefore a, b, c$ are in AP.

Hence proved.

7. Show that $x^2 + xy + y^2, z^2 + zx + x^2$ and $y^2 + yz + z^2$ are in consecutive terms of an A.P., if x, y and z are in A.P.

Solution:

x, y, z are in AP

Given, $x^2 + xy + y^2, z^2 + zx + x^2$ and $y^2 + yz + z^2$ are in AP

$$(z^2 + zx + x^2) - (x^2 + xy + y^2) = (y^2 + yz + z^2) - (z^2 + zx + x^2)$$

Let $d =$ common difference,

So, $Y = x + d$ and $x = x + 2d$

Let us consider the LHS:

$$(z^2 + zx + x^2) - (x^2 + xy + y^2)$$

$$z^2 + zx - xy - y^2$$

$$(x + 2d)^2 + (x + 2d)x - x(x + d) - (x + d)^2$$

$$x^2 + 4xd + 4d^2 + x^2 + 2xd - x^2 - xd - x^2 - 2xd - d^2$$

$$3xd + 3d^2$$

Now, let us consider RHS:

$$(y^2 + yz + z^2) - (z^2 + zx + x^2)$$

$$y^2 + yz - zx - x^2$$

$$(x + d)^2 + (x + d)(x + 2d) - (x + 2d)x - x^2$$

$$x^2 + 2dx + d^2 + x^2 + 2dx + xd + 2d^2 - x^2 - 2dx - x^2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$3xd + 3d^2$$

$$\text{LHS} = \text{RHS}$$

$\therefore x^2 + xy + y^2, z^2 + zx + x^2$ and $y^2 + yz + z^2$ are in consecutive terms of A.P

Hence proved.

EXERCISE 19.6 PAGE NO: 19.46

1. Find the A.M. between:

(i) 7 and 13 (ii) 12 and -8 (iii) $(x - y)$ and $(x + y)$

Solution:

(i) Let A be the Arithmetic mean

Then 7, A, 13 are in AP

Now, let us solve

$$A - 7 = 13 - A$$

$$2A = 13 + 7$$

$$A = 10$$

(ii) Let A be the Arithmetic mean

Then 12, A, -8 are in AP

Now, let us solve

$$A - 12 = -8 - A$$

$$2A = 12 + 8$$

$$A = 2$$

(iii) Let A be the Arithmetic mean

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Then $x - y$, A , $x + y$ are in AP

Now, let us solve

$$A - (x - y) = (x + y) - A$$

$$2A = x + y + x - y$$

$$A = x$$

2. Insert 4 A.M.s between 4 and 19.

Solution:

Let A_1, A_2, A_3, A_4 be the 4 AM Between 4 and 19

Then, 4, $A_1, A_2, A_3, A_4, 19$ are in AP.

By using the formula,

$$d = (b-a) / (n+1)$$

$$= (19 - 4) / (4 + 1)$$

$$= 15/5$$

$$= 3$$

So,

$$A_1 = a + d = 4 + 3 = 7$$

$$A_2 = A_1 + d = 7 + 3 = 10$$

$$A_3 = A_2 + d = 10 + 3 = 13$$

$$A_4 = A_3 + d = 13 + 3 = 16$$

3. Insert 7 A.M.s between 2 and 17.

Solution:

Let $A_1, A_2, A_3, A_4, A_5, A_6, A_7$ be the 7 AMs between 2 and 17

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Then, 2, A_1 , A_2 , A_3 , A_4 , A_5 , A_6 , A_7 , 17 are in AP

By using the formula,

$$a_n = a + (n - 1)d$$

$$a_n = 17, a = 2, n = 9$$

so,

$$17 = 2 + (9 - 1)d$$

$$17 = 2 + 9d - d$$

$$17 = 2 + 8d$$

$$8d = 17 - 2$$

$$8d = 15$$

$$d = 15/8$$

So,

$$A_1 = a + d = 2 + 15/8 = 31/8$$

$$A_2 = A_1 + d = 31/8 + 15/8 = 46/8$$

$$A_3 = A_2 + d = 46/8 + 15/8 = 61/8$$

$$A_4 = A_3 + d = 61/8 + 15/8 = 76/8$$

$$A_5 = A_4 + d = 76/8 + 15/8 = 91/8$$

$$A_6 = A_5 + d = 91/8 + 15/8 = 106/8$$

$$A_7 = A_6 + d = 106/8 + 15/8 = 121/8$$

\therefore the 7 AMs between 2 and 7 are $31/8$, $46/8$, $61/8$, $76/8$, $91/8$, $106/8$, $121/8$

4. Insert six A.M.s between 15 and - 13.

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Let $A_1, A_2, A_3, A_4, A_5, A_6$ be the 7 AM between 15 and -13

Then, 15, $A_1, A_2, A_3, A_4, A_5, A_6, -13$ are in AP

By using the formula,

$$a_n = a + (n - 1)d$$

$$a_n = -13, a = 15, n = 8$$

so,

$$-13 = 15 + (8 - 1)d$$

$$-13 = 15 + 7d$$

$$7d = -13 - 15$$

$$7d = -28$$

$$d = -4$$

So,

$$A_1 = a + d = 15 - 4 = 11$$

$$A_2 = A_1 + d = 11 - 4 = 7$$

$$A_3 = A_2 + d = 7 - 4 = 3$$

$$A_4 = A_3 + d = 3 - 4 = -1$$

$$A_5 = A_4 + d = -1 - 4 = -5$$

$$A_6 = A_5 + d = -5 - 4 = -9$$

5. There are n A.M.s between 3 and 17. The ratio of the last mean to the first mean is 3: 1. Find the value of n .

Solution:

Let the series be 3, $A_1, A_2, A_3, \dots, A_n, 17$

Given, $a_n/a_1 = 3/1$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

We know total terms in AP are $n + 2$

So, 17 is the $(n + 2)$ th term

By using the formula,

$$A_n = a + (n - 1)d$$

$$A_n = 17, a = 3$$

$$\text{So, } 17 = 3 + (n + 2 - 1)d$$

$$17 = 3 + (n + 1)d$$

$$17 - 3 = (n + 1)d$$

$$14 = (n + 1)d$$

$$d = 14/(n+1)$$

Now,

$$A_n = 3 + 14/(n+1) = (17n + 3) / (n+1)$$

$$A_1 = 3 + d = (3n+17)/(n+1)$$

Since,

$$a_n/a_1 = 3/1$$

$$(17n + 3) / (3n+17) = 3/1$$

$$17n + 3 = 3(3n + 17)$$

$$17n + 3 = 9n + 51$$

$$17n - 9n = 51 - 3$$

$$8n = 48$$

$$n = 48/8$$

$$= 6$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

∴ There are 6 terms in the AP

6. Insert A.M.s between 7 and 71 in such a way that the 5th A.M. is 27. Find the number of A.M.s.

Solution:

Let the series be 7, A_1 , A_2 , A_3 ,, A_n , 71

We know total terms in AP are $n + 2$

So 71 is the $(n + 2)$ th term

By using the formula,

$$A_n = a + (n - 1)d$$

$$A_n = 71, n = 6$$

$$A_6 = a + (6 - 1)d$$

$$a + 5d = 27 \text{ (5th term)}$$

$$d = 4$$

so,

$$71 = (n + 2)\text{th term}$$

$$71 = a + (n + 2 - 1)d$$

$$71 = 7 + n(4)$$

$$n = 15$$

∴ There are 15 terms in AP

7. If n A.M.s are inserted between two numbers, prove that the sum of the means equidistant from the beginning and the end is constant.

Solution:

Let a and b be the first and last terms

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

The series be $a, A_1, A_2, A_3, \dots, A_n, b$

We know, Mean = $(a+b)/2$

Mean of A_1 and $A_n = (A_1 + A_n)/2$

$$A_1 = a+d$$

$$A_n = a - d$$

$$\text{So, AM} = (a+d+b-d)/2$$

$$= (a+b)/2$$

$$\text{AM between } A_2 \text{ and } A_{n-1} = (a+2d+b-2d)/2$$

$$= (a+b)/2$$

Similarly, $(a + b)/2$ is constant for all such numbers

$$\text{Hence, AM} = (a + b)/2$$

8. If x, y, z are in A.P. and A_1 is the A.M. of x and y , and A_2 is the A.M. of y and z , then prove that the A.M. of A_1 and A_2 is y .

Solution:

Given that,

$$A_1 = \text{AM of } x \text{ and } y$$

$$\text{And } A_2 = \text{AM of } y \text{ and } z$$

$$\text{So, } A_1 = (x+y)/2$$

$$A_2 = (y+z)/2$$

$$\text{AM of } A_1 \text{ and } A_2 = (A_1 + A_2)/2$$

$$= [(x+y)/2 + (y+z)/2]/2$$

$$= [x+y+y+z]/2$$

$$= [x+2y+z]/2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Since x, y, z are in AP, $y = (x+z)/2$

$$AM = [(x + z/2) + (2y/2)]/2$$

$$= (y + y)/2$$

$$= 2y/2$$

$$= y$$

Hence proved.

9. Insert five numbers between 8 and 26 such that the resulting sequence is an A.P

Solution:

Let A_1, A_2, A_3, A_4, A_5 be the 5 numbers between 8 and 26

Then, 8, $A_1, A_2, A_3, A_4, A_5, 26$ are in AP

By using the formula,

$$A_n = a + (n - 1)d$$

$$A_n = 26, a = 8, n = 7$$

$$26 = 8 + (7 - 1)d$$

$$26 = 8 + 6d$$

$$6d = 26 - 8$$

$$6d = 18$$

$$d = 18/6$$

$$= 3$$

So,

$$A_1 = a + d = 8 + 3 = 11$$

$$A_2 = A_1 + d = 11 + 3 = 14$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$A_3 = A_2 + d = 14 + 3 = 17$$

$$A_4 = A_3 + d = 17 + 3 = 20$$

$$A_5 = A_4 + d = 20 + 3 = 23$$

EXERCISE 19.7 PAGE NO: 19.48

1. A man saved ₹ 16500 in ten years. In each year after the first he saved ₹ 100/- more than he did in the preceding year. How much did he saved in the first year?

Solution:

Given: A man saved ₹16500 in ten years

Let ₹ x be his savings in the first year

His savings increased by ₹ 100 every year.

So,

A.P will be x, 100 + x, 200 + x.....

Where, x is first term and

Common difference, $d = 100 + x - x = 100$

We know, S_n is the sum of n terms of an A.P

By using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

where, a is first term, d is common difference and n is number of terms in an A.P.

Given:

$$S_n = 16500 \text{ and } n = 10$$

$$S_{10} = 10/2 [2x + (10 - 1)100]$$

$$16500 = 5\{2x + 9(100)\}$$

$$16500 = 5(2x + 900)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$16500 = 10x + 4500$$

$$-10x = 4500 - 16500$$

$$-10x = -12000$$

$$x = 12000/10$$

$$= 1200$$

Hence, his saving in the first year is ₹ 1200.

2. A man saves ₹ 32 during the first year, ₹ 36 in the second year and in this way he increases his savings by ₹ 4 every year. Find in what time his saving will be ₹ 200.

Solution:

Given:

First year savings is ₹ 32

Second year savings is ₹ 36

In this process he increases his savings by ₹ 4 every year

Then,

A.P. will be 32, 36, 40,.....

Where, 32 is first term and common difference, $d = 36 - 32 = 4$

We know, S_n is the sum of n terms of an A.P

By using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

where, a is first term, d is common difference and n is number of terms in an A.P.

Given:

$$S_n = 200, a = 32, d = 4$$

$$S_n = n/2 [2a + (n - 1)d]$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$200 = n/2 [2(32) + (n-1)4]$$

$$200 = n/2 [64 + 4n - 4]$$

$$400 = n [60 + 4n]$$

$$400 = 4n [15 + n]$$

$$400/4 = n [15 + n]$$

$$100 = 15n + n^2$$

$$n^2 + 15n - 100 = 0$$

$$n^2 + 20n - 5n - 100 = 0$$

$$n(n + 20) - 5(n + 20) = 0$$

$$(n + 20) - 5(n + 20) = 0$$

$$(n + 20)(n - 5) = 0$$

$$n = -20 \text{ or } 5$$

$$n = 5 \text{ [Since, } n \text{ is a positive integer]}$$

Hence, the man requires 5 days to save ₹ 200

3. A man arranges to pay off a debt of ₹ 3600 by 40 annual instalments which form an arithmetic series. When 30 of the instalments are paid, he dies leaving one-third of the debt unpaid, find the value of the instalment.

Solution:

Given:

40 annual instalments which form an arithmetic series.

Let the first instalment be 'a'

$$S_{40} = 3600, n = 40$$

By using the formula,

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$S_n = n/2 [2a + (n - 1)d]$$

$$3600 = 40/2 [2a + (40-1)d]$$

$$3600 = 20 [2a + 39d]$$

$$3600/20 = 2a + 39d$$

$$2a + 39d - 180 = 0 \dots (i)$$

Given:

Sum of first 30 terms is paid and one third of debt is unpaid.

$$\text{So, paid amount} = 2/3 \times 3600 = ₹ 2400$$

$$S_n = 2400, n = 30$$

By using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

$$2400 = 30/2 [2a + (30-1)d]$$

$$2400 = 15 [2a + 29d]$$

$$2400/15 = 2a + 29d$$

$$2a + 29d - 160 = 0 \dots (ii)$$

Now, let us solve equation (i) and (ii) by substitution method, we get

$$2a + 39d = 180$$

$$2a = 180 - 39d \dots (iii)$$

Substitute the value of 2a in equation (ii)

$$2a + 29d - 160 = 0$$

$$180 - 39d + 29d - 160 = 0$$

$$20 - 10d = 0$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$10d = 20$$

$$d = 20/10$$

$$= 2$$

Substitute the value of d in equation (iii)

$$2a = 180 - 39d$$

$$2a = 180 - 39(2)$$

$$2a = 180 - 78$$

$$2a = 102$$

$$a = 102/2$$

$$= 51$$

Hence, value of first installment 'a' is ₹ 51

4. A manufacturer of the radio sets produced 600 units in the third year and 700 units in the seventh year. Assuming that the product increases uniformly by a fixed number every year, find

(i) the production in the first year

(ii) the total product in the 7 years and

(iii) the product in the 10th year.

Solution:

Given:

600 and 700 radio sets units are produced in third and seventh year respectively.

$$a_3 = 600 \text{ and } a_7 = 700$$

(i) The production in the first year

We need to find the production in the first year.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

Let first year production be 'a'

So the AP formed is, a, a+x, a+2x,

By using the formula,

$$a_n = a + (n-1)d$$

$$a_3 = a + (3-1)d$$

$$600 = a + 2d \dots (i)$$

$$a_7 = a + (7-1)d$$

$$700 = a + 6d$$

$$a = 700 - 6d \dots (ii)$$

Substitute value of a in (i) we get,

$$600 = a + 2d$$

$$600 = 700 - 6d + 2d$$

$$700 - 600 = 4d$$

$$100 = 4d$$

$$d = 100/4$$

$$= 25$$

Now substitute value of d in (ii) we get,

$$a = 700 - 6d$$

$$= 700 - 6(25)$$

$$= 700 - 150$$

$$= 550$$

∴ The production in the first year, 'a' is 550

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

(ii) the total product in the 7 years

We need to find the total product in 7 years i.e. is S_7

By using the formula,

$$S_n = n/2 [2a + (n-1)d]$$

$$n = 7, a = 550, d = 25$$

$$S_7 = 7/2 [2(550) + (7-1)25]$$

$$= 7/2 [1100 + 150]$$

$$= 7/2 [1250]$$

$$= 7 [625]$$

$$= 4375$$

∴ The total product in the 7 years is 4375.

(iii) the product in the 10th year.

We need to find the product in the 10th year i.e. a_{10}

By using the formula,

$$a_n = a + (n-1)d$$

$$n = 10, a = 550, d = 25$$

$$a_{10} = 550 + (10-1)25$$

$$= 550 + (9)25$$

$$= 550 + 225$$

$$= 775$$

∴ The product in the 10th year is 775.

5. There are 25 trees at equal distances of 5 meters in a line with a well, the distance of well from the nearest tree being 10 meters. A gardener waters all the trees separately

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

starting from the well and he returns to the well after watering each tree to get water for the next. Find the total distance the gardener will cover in order to water all the trees.

Solution:

Given: total trees are 25 and equal distance between two adjacent trees are 5 meters

We need to find the total distance the gardener will cover.

As gardener is coming back to well after watering every tree:

Distance covered by gardener to water 1st tree and return back to the initial position is $10m + 10m = 20m$

Now, distance between adjacent trees is 5m.

Distance covered by him to water 2nd tree and return back to the initial position is $15m + 15m = 30m$

Distance covered by the gardener to water 3rd tree return back to the initial position is $20m + 20m = 40m$

Hence distance covered by the gardener to water the trees are in A.P

A.P. is 20, 30, 40upto 25 terms

Here, first term, $a = 20$, common difference, $d = 30 - 20 = 10$, $n = 25$

We need to find S_{25} which will be the total distance covered by gardener to water 25 trees.

So by using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

$$S_{25} = 25/2 [2(20) + (25-1)10]$$

$$= 25/2 [40 + (24)10]$$

$$= 25/2 [40 + 240]$$

$$= 25/2 [280]$$

$$= 25 [140]$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

= 3500

∴ The total distance covered by gardener to water trees all 25 trees is 3500m.

6. A man is employed to count ₹ 10710. He counts at the rate of ₹ 180 per minute for half an hour. After this he counts at the rate of ₹ 3 less every minute than the preceding minute. Find the time taken by him to count the entire amount.

Solution:

Given: Amount to be counted is ₹ 10710

We need to find time taken by man to count the entire amount.

He counts at the rate of ₹ 180 per minute for half an hour or 30 minutes.

So, Amount to be counted in an hour = $180 \times 30 = ₹ 5400$

Amount left = $10710 - 5400 = ₹ 5310$

$S_n = 5310$

After an hour, rate of counting is decreasing at ₹ 3 per minute. This rate will form an A.P.

A.P. is 177, 174, 171,.....

Here $a = 177$ and $d = 174 - 177 = -3$

By using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

$$5310 = n/2 [2(177) + (n-1)(-3)]$$

$$5310 = n/2 [354 - 3n + 3]$$

$$5310 \times 2 = n [357 - 3n]$$

$$10620 = 357n - 3n^2$$

$$10620 = 3n(119 - n)$$

$$10620/3 = n(119 - n)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$3540 = 119n - n^2$$

$$n^2 - 119n + 3540 = 0$$

$$n^2 - 59n - 60n + 3540 = 0$$

$$n(n - 59) - 60(n - 59) = 0$$

$$(n - 59)(n - 60) = 0$$

$$n = 59 \text{ or } 60$$

We shall consider value of $n = 59$. Since, at 60th min he will count ₹ 0

∴ The total time taken by him to count the entire amount = $30 + 59 = 89$ minutes.

7. A piece of equipment cost a certain factory ₹ 600,000. If it depreciates in value 15% the first, 13.5% the next year, 12% the third year, and so on. What will be its value at the end of 10 years, all percentages applying to the original cost?

Solution:

Given: A piece of equipment cost a certain factory is ₹ 600,000

We need to find the value of the equipment at the end of 10 years.

The price of equipment depreciates 15%, 13.5%, 12% in 1st, 2nd, 3rd year and so on.

So the A.P. will be 15, 13.5, 12,..... up to 10 terms

Here, $a = 15$, $d = 13.5 - 15 = -1.5$, $n = 10$

By using the formula,

$$S_n = n/2 [2a + (n - 1)d]$$

$$S_{10} = 10/2 [2(15) + (10-1)(-1.5)]$$

$$= 5 [30 + 9(-1.5)]$$

$$= 5 [30 - 13.5]$$

$$= 5 [16.5]$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$= 82.5$$

The value of equipment at the end of 10 years is = $[100 - \text{Depreciation \%}]/100 \times \text{cost}$

$$= [100 - 82.5]/100 \times 600000$$

$$= 175/100 \times 6000$$

$$= 175 \times 600$$

$$= 105000$$

\therefore The value of equipment at the end of 10 years is ₹ 105000.

8. A farmer buys a used tractor for ₹ 12000. He pays ₹ 6000 cash and agrees to pay the balance in annual instalments of ₹ 500 plus 12% interest on the unpaid amount. How much the tractor cost him?

Solution:

Given: Price of the tractor is ₹12000.

We need to find the total cost of the tractor if he buys it in installments.

Total price = ₹ 12000

Paid amount = ₹ 6000

Unpaid amount = ₹ 12000 – 6000 = ₹ 6000

He pays remaining ₹ 6000 in 'n' number of installments of ₹ 500 each.

So, $n = 6000/500 = 12$

Cost incurred by him to pay remaining 6000 is

The AP will be:

$(500 + 12\% \text{ of } 6000) + (500 + 12\% \text{ of } 5500) + \dots$ up to 12 terms

$500 \times 12 + 12\% \text{ of } (6000 + 5500 + \dots)$ up to 12 terms)

By using the formula,

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

$$S_n = n/2 [2a + (n - 1)d]$$

$$n = 12, a = 6000, d = -500$$

$$S_{12} = 500 \times 12 + 12/100 \times 12/2 [2(6000) + (12-1)(-500)]$$

$$= 6000 + 72/100 [12000 + 11(-500)]$$

$$= 6000 + 72/100 [12000 - 5500]$$

$$= 6000 + 72/100 [6500]$$

$$= 6000 + 4680$$

$$= 10680$$

$$\text{Total cost} = 6000 + 10680$$

$$= 16680$$

∴ The total cost of the tractor if he buys it in installment is ₹ 16680.



Chapterwise RD Sharma Solutions for Class 11 Maths :

- Chapter 1–Sets
- Chapter 2–Relations
- Chapter 3–Functions
- Chapter 4–Measurement of Angles
- Chapter 5–Trigonometric Functions
- Chapter 6–Graphs of Trigonometric Functions
- Chapter 7–Values of Trigonometric Functions at Sum or Difference of Angles
- Chapter 8–Transformation Formulae
- Chapter 9–Values of Trigonometric Functions at Multiples and Submultiples of an Angle
- Chapter 10–Sine and Cosine Formulae and their Applications
- Chapter 11–Trigonometric Equations
- Chapter 12–Mathematical Induction
- Chapter 13–Complex Numbers
- Chapter 14–Quadratic Equations
- Chapter 15–Linear Inequations
- Chapter 16–Permutations
- Chapter 17–Combinations
- Chapter 18–Binomial Theorem
- Chapter 19–Arithmetic Progressions
- Chapter 20–Geometric Progressions

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

- Chapter 21–Some Special Series
- Chapter 22–Brief review of Cartesian System of Rectangular Coordinates
- Chapter 23–The Straight Lines
- Chapter 24–The Circle
- Chapter 25–Parabola
- Chapter 26–Ellipse
- Chapter 27–Hyperbola
- Chapter 28–Introduction to Three Dimensional Coordinate Geometry
- Chapter 29–Limits
- Chapter 30–Derivatives
- Chapter 31–Mathematical Reasoning
- Chapter 32–Statistics
- Chapter 33–Probability

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>

About RD Sharma

RD Sharma isn't the kind of author you'd bump into at lit fests. But his bestselling books have helped many CBSE students lose their dread of maths. Sunday Times profiles the tutor turned internet star

He dreams of algorithms that would give most people nightmares. And, spends every waking hour thinking of ways to explain concepts like 'series solution of linear differential equations'. Meet Dr Ravi Dutt Sharma — mathematics teacher and author of 25 reference books — whose name evokes as much awe as the subject he teaches. And though students have used his thick tomes for the last 31 years to ace the dreaded maths exam, it's only recently that a spoof video turned the tutor into a YouTube star.

R D Sharma had a good laugh but said he shared little with his on-screen persona except for the love for maths. "I like to spend all my time thinking and writing about maths problems. I find it relaxing," he says. When he is not writing books explaining mathematical concepts for classes 6 to 12 and engineering students, Sharma is busy dispensing his duty as vice-principal and head of department of science and humanities at Delhi government's Guru Nanak Dev Institute of Technology.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-11-maths-chapter-19-arithmetic-progressions/>